

Post-Operative Instructions

T. Ryan Pflugner, MD

MEDICATIONS

•	Pain medication and other medications are sent <u>electronically</u> to the pharmacy that you selected at our
	office. Take medication per the directions on the bottle.
	Pharmacy:

- Aspirin 325mg once daily for 30 days, if not on blood thinner prior to surgery (total joint only)
- o Celebrex 200mg, if not already on NSAID (total joint surgery only)
- o Prednisone 5mg once daily for 14 days (total joint surgery only)
- o Gabapentin 300mg 3 times a day as needed (total joint surgery only)
- o Norco 10/325 every 6 hours as needed for pain
- o Zofran as needed for nausea
- For patients at increased risk of complications with total joint surgery, 7 days of oral antibiotics will be prescribed.
- Be sure to drink plenty of water and eat small, more frequent meals to prevent constipation. You can take an over-the-counter stool softener such as Colace or MiraLAX daily, if needed.
- Restart all home medications the day after surgery, unless specified otherwise by Dr. Pflugner.
- If you had joint replacement surgery, we will manage antibiotic medication prior to having dental work. Please call (334) 793-2663 and choose the option to leave a message for the nurse. Please allow 48 business hours to process.

CARE OF OPERATIVE SITE

- Elevate the operative area when at rest. This helps to prevent excessive swelling.
- Ice can be applied to your surgical area for 20 minutes at the time, as often as you'd like, but we recommend a minimum of three times per day. This will help swelling and pain.
- If you have a large, bulky surgical dressing, it is to be removed after two days and covered with a clean, dry dressing. If you have a large, rectangular bandage dressing, it is to be removed after 7 days and covered with a clean, dry dressing until you are seen for your post-op appointment. The small incisions (made during non-total joint/arthroscopic procedures) can be covered with a clean, dry band-aid.
- You may shower. Pat area dry. Do not apply creams, ointments or lotions to the incision(s).
- If you experience drainage, redness, increase in swelling at the incision site, please call the office at (334) 793-2663 and choose the option to leave a message for the nurse.

PHYSICAL THERAPY

•	Therapy is set up prior to your surgery at the facility of your choosing. If you have not been contacted by
	that facility, <u>please call them.</u> Do not wait until your post-operative appointment.
	Facility:
_	We have also provided your a complete by material that your Deviced Therewas to be up fallow. Discontinuous

- We have also provided you a copy of the protocol that your Physical Therapist should follow. Please take it with you to your first session.
 - o If you are receiving PT in your home via Home Health agency, please be aware that after 2-4 weeks, you may transition to an outpatient PT facility.
 - Workman's Comp/Trimed cases please contact your adjuster/case manager for information about where your PT will take place.